

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/281581740>

Unit Groups of Some Classes of Power Four Radical Zero Commutative Completely Primary Finite Rings

Article · January 2014

DOI: 10.12988/ija.2014.4431

CITATIONS

2

READS

70

2 authors:



Maurice Owino Oduor

University of Kabianga

29 PUBLICATIONS 32 CITATIONS

SEE PROFILE



Michael Ojiema

Masinde Muliro University of Science and Technology

16 PUBLICATIONS 20 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Automorphisms of the Unit groups of Some Classes of finite Rings [View project](#)



A characterization of a New Structure constructed from almost contact 3-structure on a smooth differentiable manifold [View project](#)

Unit Groups of Some Classes of Power Four Radical Zero Commutative Completely Primary Finite Rings

Owino Maurice Oduor

Department of Mathematics and Computer Science
University of Kabianga
P.O. Box 2030–20200, Kericho, Kenya

Ojiema Michael Onyango

Department of Mathematics
Masinde Muliro University of Science and Technology
P. O. Box 190–50100, Kakamega, Kenya

Copyright © 2014 Owino Maurice Oduor and Ojiema Michael Onyango. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Let R be a completely primary finite ring and J be its Jacobson radical. A class of such rings in which $J^4 = (0)$, $J^3 \neq (0)$ has been constructed. Moreover, the structures of their groups of units have been determined for all the characteristics of R .

Mathematics Subject Classification: Primary 13M05, 16P10, 16U60; Secondary 13E10, 16N20

Keywords: Unit Groups, Completely Primary Finite Rings

1 Introduction

The background information on completely primary finite rings can be obtained from [1, 5]. The notations used in this paper are standard, see e.g [5].

The classification of finite rings, still remains open. Several authors have constructed finite rings whose Jacobson radical or group of units yield a particular structure. For instance, in [2], the author has obtained structures of unit groups of classes of completely primary finite rings in which the product of any three zero divisors is zero. It is well known that if R is a finite field, then the group of units is cyclic. In [3], Gilmer characterized all rings whose groups of units are cyclic. In this paper, we have constructed rings in which the product of any four zero divisors is zero. Moreover, their groups of units have been characterized. From the well known Raghavendran's result (see [5], Theorem 2), we notice that if $J^4 = (0)$, $J^3 \neq (0)$, then characteristic of R is p , p^2 , p^3 and p^4 .

2 Rings of characteristic p

Let $R' = GR(p^r, p)$ be the Galois ring of order p^r and characteristic p . Suppose U, V and W are finitely generated R' modules such that $\dim_{R'} U = s$, $\dim_{R'} V = t$, $\dim_{R'} W = \lambda$ and $s + t + \lambda = h$. Let $\{u_1, \dots, u_s\}$, $\{v_1, \dots, v_t\}$ and $\{w_1, \dots, w_\lambda\}$ be the generators of U, V and W respectively so that $R = R' \oplus U \oplus V \oplus W$ is an additive abelian group. Further, assume that $s = 1, t = 1, \lambda = h - 2$ so that $R = R' \oplus R'u \oplus R'v \oplus \sum_{j=1}^{h-2} R'w_j$ and $pu = pv = pw_j = 0, 1 \leq j \leq h - 2$. On R , define multiplication as follows:

$(r_0, r_1, r_2, \dots, r_h)(s_0, s_1, \dots, s_h) = (r_0s_0, r_0s_1 + r_1s_0, r_0s_2 + r_2s_0 + r_1s_1, r_0s_3 + r_3s_0 + r_1s_2 + r_2s_1, \dots, r_0s_h + r_h s_0 + r_1s_2 + r_2s_1)$. It is easy to verify that the given multiplication turns R into a commutative ring with identity $(1, 0, \dots, 0)$.

Proposition 1. *The ring constructed in this section is completely primary of characteristic p and*

$$J = R'u \oplus R'v \oplus \sum_{j=1}^{\lambda} R'w_j$$

$$J^2 = R'v \oplus \sum_{j=1}^{\lambda} R'w_j$$

$$J^3 = \sum_{j=1}^{\lambda} R'w_j$$

$$J^4 = (0)$$

Proposition 2. *see e.g [2]. Let R be a completely primary finite ring (not necessarily commutative). Then, the group of units, R^* of the ring R contains a cyclic subgroup $\langle b \rangle$ of order $p^r - 1$ and R^* is a semi direct product of $1 + J$ and $\langle b \rangle$.*

Proposition 3. *Let R be the ring constructed in this section, and J be its Jacobson radical. Then*

$$R^* \cong \mathbb{Z}_{p^{r-1}} \times \mathbb{Z}_{p^2}^r \times (\mathbb{Z}_p^r)^\lambda \text{ if } p \neq 2$$

for every prime integer p and positive integer r .

Proof. Let $\xi_1, \dots, \xi_r \in R'$ with $\xi_1 = 1$ such that $\overline{\xi_1}, \dots, \overline{\xi_r} \in R'/pR'$ form a basis for R'/pR' regarded as a vector space over its prime subfield F_p . For every prime integer p , $1 + J$ is a direct product of the following $r(h - 1)$ cyclic subgroups with their respective orders

$$\begin{aligned} & \{(1 + \xi_\nu u)^{\alpha_\nu} \mid 1 \leq \alpha_\nu \leq p^2\}, 1 \leq \nu \leq r \\ & \{(1 + \xi_\nu w_j)^{\beta_\nu} \mid 1 \leq \beta_\nu \leq p\}, 1 \leq \nu \leq r, 1 \leq j \leq \lambda \end{aligned}$$

The structure of R^* follows from the above Proposition 2. □

3 Rings of characteristic p^2

Let $R' = GR(p^{2r}, p^2)$ be the Galois ring of order p^{2r} and characteristic p^2 . Suppose U, V and W are finitely generated R' modules such that $\dim_{R'} U = s$, $\dim_{R'} V = t$, $\dim_{R'} W = \lambda$ and $s + t + \lambda = h$. Let $\{u_1, \dots, u_s\}, \{v_1, \dots, v_t\}$ and $\{w_1, \dots, w_\lambda\}$ be the generators of U, V and W respectively so that $R = R' \oplus U \oplus V \oplus W$ is an additive abelian group. Further, assume that $s = h - 1, t = 1, \lambda = 0$ so that $R = R' \oplus \sum_{j=1}^{h-1} R' u_j \oplus R' v$ where $pu_j \neq 0, p^2 u_j = 0, 1 \leq j \leq s$ and $pv = 0$. On R , define multiplication as follows:

$(r_0, r_1, r_2, \dots, r_{h-1}, \overline{r_h})(s_0, s_1, \dots, s_{h-1}, \overline{s_h}) = (r_0 s_0 + p \sum_{i,j=1}^{h-1} r_i s_j, r_0 s_1 + r_1 s_0, \dots, r_0 s_{h-1} + r_{h-1} s_0, r_0 \overline{s_h} + \overline{r_h} s_0)$ where $\overline{r_h}, \overline{s_h} \in R'/pR'$. It is easy to verify that the given multiplication turns R into a commutative ring with identity $(1, 0, \dots, 0, \overline{0})$.

Proposition 4. *The ring constructed in this section is completely primary of characteristic p^2 and*

$$\begin{aligned} J &= pR' \oplus \sum_{j=1}^s R' u_j \oplus R' v \\ J^2 &= pR' \oplus p \sum_{j=1}^s R' u_j \oplus R' v \\ J^3 &= p \sum_{j=1}^s R' u_j \\ J^4 &= (0). \end{aligned}$$

Proposition 5. *Let R be the ring constructed in this section, and J be its Jacobson radical. Then*

$$R^* \cong \mathbb{Z}_{p^{r-1}} \times (\mathbb{Z}_{p^2}^r)^s \times (\mathbb{Z}_p^r)^2$$

for every prime integer p and positive integer r .

Proof. Let $\xi_1, \dots, \xi_r \in R'$ with $\xi_1 = 1$ such that $\overline{\xi_1}, \dots, \overline{\xi_r} \in R'/pR'$ form a basis for R'/pR' regarded as a vector space over its prime subfield F_p . It therefore suffices to characterize $1 + J$, for the structure of R^* will easily follow from Proposition 2.

For every prime integer p , $1 + J$ is a direct product of the following $r(s + 2)$ cyclic subgroups with their respective orders

$$\begin{aligned} & \{(1 + p\xi_\nu)^\alpha \mid 1 \leq \alpha \leq p\}, 1 \leq \nu \leq r \\ & \{(1 + \xi_\nu u_j)^{\beta_j} \mid 1 \leq \beta_j \leq p^2\}, 1 \leq \nu \leq r, 1 \leq j \leq s \\ & \{(1 + p\xi_\nu v)^{k_\nu} \mid 1 \leq k_\nu \leq p\}, 1 \leq \nu \leq r \end{aligned}$$

The structure of R^* follows from the above Proposition 2. □

4 Rings of characteristic p^3

Let $R' = GR(p^{3r}, p^3)$ be the Galois ring of order p^{3r} and characteristic p^3 . Suppose U, V and W are finitely generated R' modules such that $\dim_{R'} U = s, \dim_{R'} V = t, \dim_{R'} W = \lambda$ and $s + t + \lambda = h$. Let $\{u_1, \dots, u_s\}, \{v_1, \dots, v_t\}$ and $\{w_1, \dots, w_\lambda\}$ be the generators of U, V and W respectively so that $R = R' \oplus U \oplus V \oplus W$ is an additive abelian group. Further, assume that $s = h - 1, t = 1, \lambda = 0$ so that $R = R' \oplus \sum_{j=1}^{h-1} R' u_j \oplus R' v$ where $p^2 u_j \neq 0, p^3 u_j = 0, 1 \leq j \leq s$ and $p v = 0$. On R , define multiplication as follows:

$(r_0, \overline{r_1}, \overline{r_2}, \dots, \overline{r_{h-1}}, \tilde{r}_h)(s_0, \overline{s_1}, \dots, \overline{s_{h-1}}, \tilde{s}_h) = (r_0 s_0, r_0 \overline{s_1} + \overline{r_1} s_0, \dots, r_0 \overline{s_{h-1}} + \overline{r_{h-1}} s_0, r_0 \tilde{s}_h + \tilde{r}_h s_0 + \sum_{i,j=1}^{h-1} \overline{r_i} \overline{s_j})$ where $\overline{r_i}, \overline{s_j} \in R'/p^2 R'$ and $\tilde{r}_h, \tilde{s}_h \in R'/pR'$. It is readily verified that the given multiplication turns R into a commutative ring with identity $(1, \overline{0}, \dots, \overline{0}, \tilde{0})$.

Proposition 6. *The ring constructed in this section is completely primary of characteristic p^3 and*

$$\begin{aligned} J &= pR' \oplus \sum_{j=1}^s R' u_j \oplus R' v \\ J^2 &= p^2 R' \oplus p \sum_{j=1}^s R' u_j \oplus R' v \end{aligned}$$

$$J^3 = pR'v$$

$$J^4 = (0).$$

Proposition 7. *Let R be the ring constructed in this section, and J be its Jacobson radical. Then its group of units is characterized as follows:*

$$R^* \cong \begin{cases} \mathbb{Z}_{2^{r-1}} \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4^{r-1} \times \mathbb{Z}_8^r \times \mathbb{Z}_4^r \times (\mathbb{Z}_2^r)^{s-1} & \text{if } p = 2 \\ \mathbb{Z}_{p^{r-1}} \times \mathbb{Z}_{p^2}^r \times \mathbb{Z}_{p^2}^r \times (\mathbb{Z}_{p^2}^r)^s & \text{if } p \neq 2 \end{cases}$$

Proof. Let $\xi_1, \dots, \xi_r \in R'$ with $\xi_1 = 1$ such that $\bar{\xi}_1, \dots, \bar{\xi}_r \in R'/pR'$ form a basis for R'/pR' regarded as a vector space over its prime subfield F_p . If $p = 2$, consider $\nu = 1, \dots, r$ and $y \in R'$ such that $x^2 + x + \bar{y} = \bar{0}$ over R'/pR' has no solution in the field R'/pR' . We easily notice that $1 + J$ is a direct product of the following $r(s + 2) + 1$ cyclic subgroups with their respective orders.

$$\begin{aligned} & \{(-1 + 4\xi_1)^\alpha \mid \alpha = 1, 2\} \\ & \{(1 + 4y)^\beta \mid \beta = 1, 2\} \\ & \{(1 + 2\xi_\nu)^{\kappa_\nu} \mid \kappa_\nu = 1, \dots, 4\}, 2 \leq \nu \leq r \\ & \{(1 + 2\xi_\nu u_j)^{\tau_{j\nu}} \mid \tau_{j\nu} = 1, 2\}, \nu = 1, \dots, r; j = 1, \dots, s - 1 \\ & \{(1 + \xi_\nu u_{s-1} + \xi_\nu u_s)^{\gamma_\nu} \mid \gamma_\nu = 1, \dots, 4\}, \nu = 1, \dots, r \\ & \{(1 + \xi_\nu u_s + \xi_\nu v)^{\delta_\nu} \mid \delta_\nu = 1, \dots, 8\}, \nu = 1, \dots, r \end{aligned}$$

If $p \neq 2$, $1 + J$ is a direct product of the following $r(s + 2)$ cyclic subgroups with their respective orders:

$$\begin{aligned} & \{(1 + p\xi_\nu)^{\alpha_\nu} \mid \alpha_\nu = 1, \dots, p^2\}, \nu = 1, \dots, r \\ & \{(1 + \xi_\nu u_j)^{\beta_{j\nu}} \mid \beta_{j\nu} = 1, \dots, p^2\}, j = 1, \dots, s; \nu = 1, \dots, r \\ & \{(1 + \xi_\nu v)^{\alpha_\nu} \mid \alpha_\nu = 1, \dots, p^2\}, \nu = 1, \dots, r. \end{aligned}$$

In both cases, the structure of R^* follows from Proposition 2. □

5 Rings of characteristic p^4

Let $R' = GR(p^{4r}, p^4)$ be the Galois ring of order p^{4r} and characteristic p^4 . Suppose U, V and W are finitely generated R' modules such that $\dim_{R'} U = s$, $\dim_{R'} V = t$, $\dim_{R'} W = \lambda$ and $s + t + \lambda = h$. Let $\{u_1, \dots, u_s\}, \{v_1, \dots, v_t\}$ and $\{w_1, \dots, w_\lambda\}$ be the generators of U, V and W respectively so that $R = R' \oplus U \oplus V \oplus W$ is an additive abelian group. Further, assume that $s = h, t = 0, \lambda = 0$ so

that $R = R' \oplus \sum_{j=1}^s R' u_j$ where $pu_j = 0$, $1 \leq j \leq s$. On R define multiplication as follows:

$(r_0, \overline{r_1}, \overline{r_2}, \dots, \overline{r_h})(s_0, \overline{s_1}, \dots, \overline{s_h}) = (r_0 s_0, r_0 \overline{s_1} + \overline{r_1} s_0, \dots, r_0 \overline{s_h} + \overline{r_h} s_0)$ where $\overline{r_i}, \overline{s_j} \in R'/pR'$, $1 \leq i, j \leq s$. This multiplication turns R into a commutative ring with identity $(1, \overline{0}, \dots, \overline{0})$.

Proposition 8. *The ring constructed in this section is completely primary of characteristic p^4 and*

$$\begin{aligned} J &= pR' \oplus \sum_{j=1}^s R' u_j \\ J^2 &= p^2 R' \\ J^3 &= p^3 R' \\ J^4 &= (0). \end{aligned}$$

Proposition 9. *Let R be the ring constructed in this section, and J be its Jacobson radical. Then its group of units is characterized as follows:*

$$R^* \cong \begin{cases} \mathbb{Z}_{2^{r-1}} \times \mathbb{Z}_2 \times \mathbb{Z}_4 \times \mathbb{Z}_8^{r-1} \times (\mathbb{Z}_2)^s & \text{if } p = 2 \\ \mathbb{Z}_{p^{r-1}} \times \mathbb{Z}_{p^3}^r \times (\mathbb{Z}_p)^s & \text{if } p \neq 2 \end{cases}$$

Proof. See Proposition 5 in [4]. □

References

- [1] Y. Alkhamees, Finite completely primary finite rings in which the product of any two zero divisors of a ring is in its coefficient subring, *IJMMS.*, **17**(3), (1994), 463-468.
- [2] C.J. Chikunji, On unit groups of completely primary finite rings, *Mathematical Journal of Okayama University*, **50**, (2008)
- [3] R.W. Gilmer, Finite rings having a cyclic multiplicative group of units, *Amer. J. Math.*, **85**, (1963), 447-452
- [4] O. M. Oduor, O. M. Onyango, E. Mmasi, Units of commutative completely primary finite rings of characteristic p^n , *International Journal of Algebra.*, **7**(6), (2013), 259-266.

- [5] R. Raghavendran, Finite associative rings, *Compositio Math.*, **21**(2), (1969),195-229.

Received: March 21, 2014